

FIFRA Section 2(ee) Recommendation

**For Control of Botrytis Leaf Blight on Onions (Crop Group 3)
FOR USE AND DISTRIBUTION IN THE STATE OF TEXAS ONLY
Expires 12/31/2026**

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the US Environmental Protection Agency.

All applicable directions, restrictions, and precautions on the EPA-registered labels must be followed. Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA §2(ee) recommendations in your respective state.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Onions (Crop Group 3)

Pathogen	Application Rate	Use Instructions
Botrytis Leaf Blight (<i>Botrytis squamosa</i>)	Apply at a concentration of 4.5 ounces (128 grams) per 100 gallons of water	Begin as a preventative spray. Repeat every 7-14 days as part of a rotational program with fungicides labelled for this use.

Crop-Specific Restrictions:

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply when weather conditions favor drift or runoff from treated areas.

Do not allow the mixture to stand overnight in the spray tank.

Refer to EPA-approved label for complete list of **USE RESTRICTIONS**.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the precautionary statements plus all other information appearing on the product labels. **This recommendation must be in the possession of the user at the time of pesticide application.**

