

FIFRA Section 2(ee) Recommendation

For suppression of Root Knot (*Meloidogyne* spp.), Cyst (*Globodera* spp.), Lesion (*Pratylenchus* spp.), Ring (*Mesocriconema* spp.) and Spiral (*Helicotylenchus* spp.) nematodes

FOR USE AND DISTRIBUTION IN THE STATE OF TEXAS ONLY
Expires 12/31/2027

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the US Environmental Protection Agency.

All applicable directions, restrictions, and precautions on the EPA-registered labels must be followed. Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA §2(ee) recommendations in your respective state.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop: All labeled crops

Application Rate: 1-2 quarts per acre as part of a root disease program for nematode suppression.

Root disease soil application: Mix the required amount in sufficient water to apply by one of the following methods:

- Soil drench at transplanting, using a “water wheel” injector, spray nozzles.hoses, or other method to drench each root ball and/or planting hole.
- Soil or seed line drench, or banded spray (in-furrow) at planting (refer to label specific instructions).

Follow-up (post planting): Preventative applications can be made every 2-4 weeks as needed by one or more of the following methods:

- Drip (trickle) or any type of sprinkler irrigation, any time after planting or transplanting. See Chemigation Instructions for more information.
- Directed spray onto the soil surface and/or lower plant parts. If targeting root disease, follow immediately with sufficient overhead sprinkler irrigation to move Double Nickel[®] LC to the root zone.
- Injection directly into the rooting zone using shanks or similar equipment.





Double Nickel® LC

BIOFUNGICIDE

EPA Reg. No. 70051-107

Crop-Specific Restrictions:

Do not apply when wind speed favors drift beyond the area intended for treatment.

Refer to EPA-approved label for complete list of **USE RESTRICTIONS**.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the precautionary statements plus all other information appearing on the product labels. **This recommendation must be in the possession of the user at the time of pesticide application.**

