

FIFRA Section 2(ee) Recommendation

For Control of *Botrytis cinerea* in grapes in NY only

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the US Environmental Protection Agency.

All applicable directions, restrictions, and precautions on the EPA-registered labels must be followed. Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA §2(ee) recommendations in your respective state.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop: Grapes

Application Rate: 4.5 oz/100 gal in sufficient water to provide uniform coverage (5 gal/A minimum for aerial application)

Apply LifeGard[®] WG at 4.5 oz/100 gallons for control of *Botrytis cinerea*. Time applications to start 3-5 days before suspected infection. Repeat every 7 to 14 days as part of a rotational program with fungicides labeled for this use.

Crop-Specific Restrictions:

Do not apply to plants that are stressed due to drought, excessive moisture, excessive hot or cold temperatures, herbicide injury, or other environmental stresses.

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply when weather conditions favor drift or runoff from treated areas.

Do not allow the mixture to stand overnight in the spray tank.

Refer to EPA-approved label for complete list of **USE RESTRICTIONS**.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the precautionary statements plus all other information appearing on the product labels. **This recommendation must be in the possession of the user at the time of pesticide application.**