

# CYD-X<sup>®</sup> HP

## INSECTICIDAL VIRUS

### High Potency Aqueous Suspension Biological Insecticide for Control of the Codling Moth

**Active Ingredient\*:**

<i>Cydia pomonella</i> granulovirus	0.06%
<b>Other Ingredients:</b>	99.94%
<b>Total</b>	100.00%

\*Contains at least  $3 \times 10^{13}$  viral occlusion bodies (OBs) per Liter.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See Inside Panels for Additional Precautionary Statements, Directions for Use, and Storage and Disposal

CYD-X<sup>®</sup> HP is an insecticidal virus product for control of larvae of the codling moth. Read this label carefully. Refer to the Technical Bulletin for Additional Guidelines for Use.

**Net Contents:**

Lot No.:  
EPA Reg. No. 70051-112  
EPA Est. No.

Manufactured by  
Certis USA LLC  
9145 Guilford Road  
Suite 175  
Columbia, MD 21046

**CERTIS**

ESL20240611  
ESN20240904  
REV20250624



FOR ORGANIC PRODUCTION

**FIRST AID**

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**Hotline Number**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. 1-800-255-3924 (24-hour emergency response).

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS (AND DOMESTIC ANIMALS):****CAUTION:**

Harmful if inhaled. Avoid breathing spray mist. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

**PERSONAL PROTECTIVE EQUIPMENT:**

Applicators and other handlers must wear:

- Long sleeved shirt and long pants
- Shoes plus socks
- Protective eyewear such as goggles, face shield, or shielded safety glasses.

Mixers/loaders and applicators must wear a minimum of dust/mist filtering respirator meeting NIOSH standards of at least N-95, R-95, or P-95. Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Control Statements:**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [(40 CFR 170.607 (d) and (e) (f)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**IMPORTANT:** When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

**USER SAFETY RECOMMENDATION****Users Should:**

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS:**

For terrestrial uses. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

*This is a Specimen Label. It may not reflect the most-recent approved label for use in your state. Always refer to the label on the product packaging for approved use instructions. Please contact your Certis sales representative for more information.*

## GENERAL INFORMATION

### CYD-X® HP BIOLOGICAL INSECTICIDE

#### PRODUCT DESCRIPTION

CYD-X® HP contains an insecticidal baculovirus (scientific name: *Cydia pomonella* granulovirus or CpGV) that is specific to larvae of the codling moth (*Cydia pomonella*). Because of this specificity to the codling moth, CYD-X® HP provides a unique opportunity for biological control of codling moth without harmful effects on humans, domestic animals, wildlife, beneficial insects and plants.

#### BIOLOGICAL PROPERTIES and MODE OF ACTION

CYD-X® HP is a biological insecticide which must be ingested by larvae in order to be effective. Upon ingestion, the viral occlusion bodies (OBs) dissolve in the larval midgut and release infectious virions. These enter the cells lining the digestive tract and replicate in the nuclei of those cells. The resulting progeny virus infects other tissues within the larva, which stops feeding and eventually dies from massive viral infection. After death, the larva disintegrates, releasing billions of new OBs which may infect other codling moth larvae upon ingestion. Death occurs 3 to 7 days after ingestion of CYD-X® HP, depending on dosage and ambient temperature. Death may occur more quickly at higher temperatures and higher dosage.

Baculoviruses have been found only in invertebrates; currently there are no reports of members of this family infecting vertebrates or plants. CYD-X® HP virions do not reproduce in mammalian cells.

Bioactivity of CYD-X® HP may decrease if exposed to temperatures above 86°F for more than 3 months. For best results, keep the product refrigerated or frozen when not in use.

## MIXING INSTRUCTIONS

**Mix CYD-X® HP with water and apply as a spray to fruit and foliage. For best results, use water at or near neutral pH (7).**

#### APPLICATION RATES

CROP	Application rate
Apple, pear, plum, prune, and walnut	1/2 to 3 fluid ounces per acre of trees
<b>Comments</b> <p>Apply in sufficient water to thoroughly wet the tree canopy without excessive runoff. Select a spray volume appropriate to tree size and density of foliage. For small trees use 50-100 gallons of spray per acre. Larger trees with denser canopy may require 100-200 gallons per acre. Consult your local Cooperative Extension Service if you need advice about how to attain good spray coverage in your orchard.</p> <p>Make at least two (2) applications of CYD-X® HP per codling moth larval generation. Target small larvae early in their life cycle. Repeat as necessary to maintain control. CYD-X® HP breaks down under direct exposure to sunlight. If codling moths are still flying and laying eggs, reapply after 7-8 sunny days.</p> <p>If using temperature models (heat unit accumulation measured in degree days) and pheromone trap catch to predict codling moth development, make the first spray against the first larval generation around 250 degree days after codling moth Biofix (first moth flight). If targeting the second generation, start at approx. 1100-1200 degree days after biofix. "No Biofix" models (using heat units only, with no trap catch information) are now used in some growing regions. Consult your local Cooperative Extension Service for information about using degree day models to guide spray timing.</p> <p>If not using heat units to time sprays, apply CYD-X® HP in the 1st and 2nd cover sprays against first generation larvae. For second generation, apply in the 4th and 5th cover sprays.</p> <p>Frequent applications at low rates may be more effective than one or two applications at high rate.</p>	

CYD-X® HP can be applied up to and including the day of harvest and storage.

## STORAGE AND DISPOSAL

#### PESTICIDE STORAGE:

Store this product in original sealed container in a cool, dry place inaccessible to children and pets. Store at temperatures below 86°F. Bioactivity may be impaired by storage above 86°F. Storing the product in a freezer or refrigerator will extend its shelf life.

#### PESTICIDE DISPOSAL:

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

#### CONTAINER HANDLING:

Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if appropriate. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by State and local ordinances.

#### WARRANTY

To the extent consistent with applicable law, Certis USA LLC warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the insect problem, condition of the crop, incompatibility with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herein. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OR MERCHANTABILITY IS MADE.**