

## FIFRA Section 2(ee) Recommendation

### For Control of *Botrytis cinerea*, *Neofabraea malicorticis*, and *Phaciopycnis washingtonensis* in Apples and Pears in OR and WA only

***This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the US Environmental Protection Agency.***

**All applicable directions, restrictions, and precautions on the EPA-registered labels must be followed.** Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA §2(ee) recommendations in your respective state.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### ***Crop: Apples, Pears***

DISEASE	APPLICATION RATE	MAXIMUM ANNUAL RATE	SPECIFIC USE INSTRUCTIONS
<i>Botrytis cinerea</i> <i>Neofabraea malicorticis</i> <i>Phaciopycnis washingtonensis</i>	0.5 to 1.0 gal. of product mixed with 30 to 100 gal. of water	102 gal/acre (16 lbs. Cu/acre)	<p>Apply using standard ground spray equipment. If using ultra-low volume sprayers, mix 0.5 to 1.0 gal. in 5 to 30 gal. of water and apply to one acre. For application by aircraft, mix 0.5 to 1.0 gal. of product with 5-40 gallons of water and apply to one acre.</p> <p>Begin treatment when disease first appears and reapply at 7 to 10 day intervals for as long as needed. Use the higher rate following heavy rain or when the amount of disease is increasing rapidly. If possible, time applications so that 12 hours of dry weather follow application. Use the higher rate to control diseases that may go dormant and overwinter.</p> <p>As a dormant or delayed dormant application, up to 200 gallons diluted spray/acre may be applied. For tender plants test for phytotoxicity before treating large areas.</p>
	6.4 to 12.8 fl. oz. of product with 10 gallons of water.	299 fl. oz./1000 ft <sup>2</sup> (16 lbs. Cu/acre)	<p>Apply 1.1 to 2.3 gallons of diluted spray using standard ground spray equipment to 1,000 ft<sup>2</sup>.</p> <p>Begin treatment when disease first appears and reapply at 7 to 10 day intervals for as long as needed. Use the higher rate following heavy rain or when the amount of disease is increasing rapidly. If possible, time applications so that 12 hours of dry weather follow application. Use the higher rate to control diseases that may go dormant and overwinter.</p> <p>For tender plants test for phytotoxicity before treating large areas.</p>





## Cueva® Fungicide Concentrate

Flowable Liquid Copper Fungicide

EPA Reg. No. 67702-2-70051

### Crop-Specific Restrictions:

- Do not exceed one application during the fall, late dormant period.
- Do not exceed one application between silver tip and green tip growth stages.
- Do not reapply within 5 days during the bloom and growing stages.
- Do not apply more than 6 lbs. Cu/acre.
- May cause russetting of susceptible apple varieties.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the precautionary statements plus all other information appearing on the product labels. **For Washington only, this recommendation is not required to be in the possession of the user at the time of pesticide application. For all listed states other than Washington, this recommendation must be in the possession of the user at the time of pesticide application.**

