

**FIFRA Section 2(ee) Recommendation****For Control of *Alternaria alternata* on Almonds and Pistachios  
FOR USE AND DISTRIBUTION IN THE STATE OF CALIFORNIA ONLY**

***This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the US Environmental Protection Agency.***

**All applicable directions, restrictions, and precautions on the EPA-registered labels must be followed.** Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA §2(ee) recommendations in your respective state.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

***Crops: Almond & Pistachio*****Application Rate: 1-2 quarts per acre**

Apply Double Nickel<sup>®</sup> LC when conditions are conducive for development of disease. Repeat applications every 7-21 days as needed, for as long as conditions favor development of disease. Thorough spray coverage of plant foliage will result in best performance. For improved control, mix or rotate Double Nickel<sup>®</sup> LC with other fungicides approved for such use.

**Crop-Specific Restrictions:**

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply when weather conditions favor drift or runoff from treated areas.

Refer to EPA-approved label for complete list of **USE RESTRICTIONS**.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the precautionary statements plus all other information appearing on the product labels. **This recommendation must be in the possession of the user at the time of pesticide application.**

