

FIFRA Section 2(ee) Recommendation
For Control of Tar Spot on Corn (Crop Group 15)
FOR USE AND DISTRIBUTION IN THE STATE OF TEXAS ONLY
Expires 12/31/2026

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the US Environmental Protection Agency.

All applicable directions, restrictions, and precautions on the EPA-registered labels must be followed. Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA §2(ee) recommendations in your respective state.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Corn (Crop Group 15)

Target Disease	Use Rate	Additional Information
Tar Spot (<i>Phyllachora maydis</i>)	Apply product at 0.5-1 qt/A in sufficient water to achieve thorough coverage of the crop with minimal runoff.	Apply preventatively prior to disease onset as a foliar spray. Repeat applications as needed within 3-to-10-day intervals if conditions favor disease development and use as part of an integrated program.

Crop-Specific Restrictions:

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply when weather conditions favor drift or runoff from treated areas.

TO ENSURE UNIFORMITY, MIX PRODUCT WELL BEFORE USE.

Refer to EPA-approved label for complete list of **USE RESTRICTIONS**.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the precautionary statements plus all other information appearing on the product labels. **This recommendation must be in the possession of the user at the time of pesticide application.**

