# **OSO**° 5% SC

FUNGICIDE

## STRIKE OUT FUNGAL DISEASES WITH THE POWER OF OSO<sup>®</sup> 5% SC FUNGICIDE

Preventative or curative treatment of fungal diseases, including Botrytis, Alternaria, anthracnose, powdery mildew, and leaf and fruit rots in fruit and vegetable crops.



## **PRODUCT HIGHLIGHTS**

With a novel mode of action, polyoxin D zinc salt is the only active ingredient in the FRAC group 19, making it an ideal choice for resistance management

Can be used as a standalone or easily integrated into existing IPM programs

Comes in an easy-to-apply suspension concentrate (SC) liquid formulation, and is labeled for ground, aerial, and chemigation

OSO® 5% SC is approved for use in organic production (OMRI-Listed® WSDA -Listed®, and NOP-Compliant) and has efficacy comparable to many conventional products

A residue-exempt material with a 4hr REI and a 0 day PHI







1.800.250.5024 www.CertisBio.com

#### **HOW IT WORKS**

The active ingredient of OSO® 5% SC is polyoxin D zinc salt derived from a natural fermentation process. Polyoxin D zinc salt is a chitin synthase inhibitor that controls plant pathogenic fungi by inhibiting the formation of chitin, a vital component of fungal cell walls. When the pathogen comes in contact with OSO, early growth or colonization is stopped and cannot spread to new plant tissue.

#### **ONLY ACTIVE INGREDIENT IN FRAC GROUP 19**

#### **EFFECTS OF POLYOXIN-D ON** FUNGAL SPORE GERMINATION

#### EFFECTS OF POLYOXIN-D ON **CELL WALL INTEGRITY**



### **HOW TO APPLY**

- Shake well before use, fill tank with water to ½ of the intended final volume.
- Start agitation of the spray tank, then add the appropriate amount of product to the tank according to the rates in the label.
- Agitate to ensure thorough mixing while adding the remaining required water.
- Do not allow the mixture to stand without agitation.
- Mix only the amount of solution needed to treat the desired area.

#### **TANK MIXING**

#### TIMING

See product label

for application

timing based on

crops and diseases.

When tank mixing OSO 5% SC with other products, observe all precautions and limitations on each separate product label. When planning to mix this product with others, it is advisable to conduct a "jar test" to determine the physical compatibility of this product with the others.

#### **RECOMMENDED RATES**

6.5 – 13oz/acre - Use the 6.5oz/acre for preventative applications when disease is expected but not visible, in periods of low disease pressure, or in a tank mix with other fungicides for resistance management. A higher rate should be used when disease pressure is high or in stand-alone applications.

#### **USE WITH OTHER TREATMENTS**

If more applications or shorter intervals than indicated on the label are needed to maintain disease control, alternate OSO with other fungicides having different modes of action to prevent or slow development of pathogen resistance.

#### Storage

Store in dry place away from food or feed.

**PHI/REI** 

4 hour REI; 0 day PHI



#### **ACTIVE INGREDIENT** Polyoxin D zinc salt 5.0%

Please see label for complete list of labeled crops and diseases

#### **DISEASES CONTROLLED**

**Cucurbits**: Anthracnose **Powdery Mildew** Early Blight Scab Gray Mold **Target Leaf Spot Gummy Stem Blight** 

**Pome and Stone Fruits:** Alternaria leaf spot Leaf Blotch **Powdery Mildew** 

Scah Botrytis Blossom Blight

Potatoes: Black Scurf **Early and Late** Blight

White Mold

Citrus: Alternaria Brown Spot **Botrytis Rot** 

Septoria Spot

#### Grapes

Alternaria Leaf Spot Anthracnose Gray Mold

**Powdery Mildew Yellow Rust** 

#### MAIN CROPS

Root and Tubers Vegetables Bulb Vegetables Brassica (Cole) Leafy Vegetables Legumes **Fruiting Vegetables** Cucurbits

**Citrus Fruits Pome Fruits Stone Fruits** Berries and Small Fruits Herbs and Spices



1.800.250.5024 www.CertisBio.com

Copyright 2021 Certis USA LLC.