

FIFRA Section 2(ee) Recommendation
For Use on Tobacco to Control Angular Leaf Spot (*Pseudomonas syringae* pv. *tabaci*) in FL, GA, KY, NC, SC, and VA only

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the US Environmental Protection Agency.

All applicable directions, restrictions, and precautions on the EPA-registered labels must be followed. Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA §2(ee) recommendations in your respective state.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop: Tobacco

DISEASE	TYPE OF APPLICATION	APPLICATION RATE	MAXIMUM ANNUAL RATE	SPECIFIC USE INSTRUCTIONS
<i>Angular Leaf Spot (Pseudomonas syringae pv. tabaci)</i>	Standard Ground Spray Equipment (Except Ultra Low Volume Sprayers)	Mix 0.5 to 2.0 gal. of product with 30 to 100 gal. of water / acre	25 gal/acre (4 lbs. Cu/acre)	Begin treatment when disease first appears and reapply at 10-day intervals for as long as needed. Use the higher rate following heavy rain or when the amount of disease is increasing rapidly. If possible, time applications so that 12 hours of dry weather follow application. Use the higher rate to control diseases that may go dormant and overwinter. This product may cause some copper toxicity on some plant species. For tender plants, test for phytotoxicity before treating large areas.
		For smaller sizes: Mix 6.4 to 25.6 fl. oz. of product with 10 gal. of water. Apply 1.1 to 2.3 gal. of diluted spray / 1,000 ft ² .	73.5 fl. oz./1000 ft ² (4 lbs. Cu/acre)	
	Ultra Low Volume Sprayers	Mix 0.5 to 2.0 gal. of product in 5 to 30 gal. of water / acre	25 gal/acre (4 lbs. Cu/acre)	
		For smaller sizes: Mix 1.5 to 6 fl. oz. of this product with 15 to 88 fl. oz. water and apply to 1000 sq. ft.	73.5 fl. oz./1000 ft ² (4 lbs. Cu/acre)	
Aircraft Applications	Mix 0.5 to 2.0 gal. of product with 5-40 gallons of water / acre	25 gal/acre (4 lbs. Cu/acre)		





Cueva® Fungicide Concentrate

Flowable Liquid Copper Fungicide

EPA Reg. No. 67702-2-70051

Crop-Specific Restrictions:

Use on tobacco in transplant beds (or on field grown plants).
Do not reapply within 10 days.
Do not apply more than 2 lbs. Cu/acre.

Refer to EPA-approved label for complete list of **USE RESTRICTIONS**.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the precautionary statements plus all other information appearing on the product labels. **This recommendation must be in the possession of the user at the time of pesticide application.**

