

FIFRA Section 2(ee) Recommendation**For control of *Cercospora* leaf spot on beets in CA, CO, IL, MI, and WI**

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the US Environmental Protection Agency.

All applicable directions, restrictions, and precautions on the EPA-registered labels must be followed. Please contact your regional sales manager to determine the specific requirements for FIFRA §2(ee) recommendations in your respective state.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crops: Beets**Application Rate: 1-2 quarts per acre**

Apply Double Nickel® LC when conditions are conducive for development of disease. Repeat applications every 3-10 days as needed, for as long as conditions favor development of disease. Thorough spray coverage of plant foliage will result in best performance. Mix or rotate Double Nickel® LC with other fungicides, such as Cueva®, for improved performance.

Crop-Specific Restrictions:

Do not apply when wind speed favors drift beyond the area intended for treatment.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the precautionary statements plus all other information appearing on the product labels. **This recommendation must be in the possession of the user at the time of pesticide application.**

Double Nickel is a registered trademark of Certis USA LLC.
Cueva is a registered trademark of W. Neudorff GmbH KG.

