

FIFRA Section 2(ee) Recommendation

For control of algal stem blotch (*Cephaleurus virescens*) on blueberry in AL, FL, GA, NC, and SC only

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the US Environmental Protection Agency.

All applicable directions, restrictions, and precautions on the EPA-registered labels must be followed. Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA §2(ee) recommendations in your respective state.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop: Blueberry

Application Rate: 1.75 – 3.5 lb per acre

Apply Kocide® 3000-O at the first sight of algal stem blotch. Thorough spray coverage of plant foliage will result in best performance. Start applications after harvest on a monthly interval.

Crop-Specific Restrictions:

Do not apply when wind speed favors drift beyond area intended for treatment.

If Kocide® 3000-O is applied in a spray solution having a pH of less than 6.5, phytotoxicity may occur.

Do not tank mix Kocide® 3000-O with “Aliette” fungicide for use on any registered crops unless appropriate precautions have been taken to buffer the spray solution because severe phytotoxicity may result.

Refer to EPA-approved label for complete list of **USE RESTRICTIONS**.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the precautionary statements plus all other information appearing on the product labels. **This recommendation must be in the possession of the user at the time of pesticide application.**

