

**FIFRA Section 2(ee) Recommendation**  
**For Control of Melanose, Scab, Greasy Spot, Pink Pitting, Brown Rot, and Septoria Spot in Citrus in Florida and Arizona**

***This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the US Environmental Protection Agency.***

**All applicable directions, restrictions, and precautions on the EPA-registered labels must be followed.** Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA §2(ee) recommendations in your respective state.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Crop: Citrus**

<b>CITRUS</b>			
<b>Grapefruit, Kumquat, Lemon, Lime, Orange, Pummelo, Tangelo and Tangerine</b>			
<p>Kocide® 50DF may be mixed with dry foliar nutritionals (micronutrients) to create “Shot Bag” mixes to meet the various nutritional requirements of citrus and provide disease protection as described on this label. Kocide® 50DF per acre rates in these mixes must not exceed the maximum labeled rates for disease control.</p> <p>Adding foliar nutritionals or other products to spray mixtures containing Kocide® 50DF and applying to citrus during the post-bloom period when young fruit are present may result in spray burn.</p>			
<b>Disease</b>	<b>Application Rate/Acre</b>	<b>Maximum Annual Rate/Acre</b>	<b>Use Instructions</b>
Melanose, Scab	2 – 6.3 lbs. (1 - 3.15 lbs. <i>metallic copper</i> )	25.1 lbs. (12.6 lbs. <i>metallic copper</i> )	Apply as pre-bloom and post-bloom sprays. Use the higher rates when conditions favor disease. Minimum retreatment interval is 7 days.
Greasy Spot, Pink Pitting	2 – 6.3 lbs. (1 - 3.15 lbs. <i>metallic copper</i> )		Apply in summer on expanded new flush. Repeat on subsequent flushes where disease pressure is severe. Use the higher rates when conditions favor disease. Minimum retreatment interval is 7 days.
Brown Rot, Septoria Spot	2 – 6.3 lbs. (1 - 3.15 lbs. <i>metallic copper</i> )		Begin application in fall before or just after the first rain and continue if needed. For Brown Rot only, apply to skirts of trees to a height of at least 4 feet. For control of Septoria Spot or where fruit have already been infected with Brown Rot, apply to entire tree. Apply also to bare ground one foot beyond skirt.



			Use the higher rates when conditions favor disease. Minimum retreatment interval is 7 days.
<b>NOTE:</b> Phytotoxicity may occur on young tender flush when Kocide® 50DF is applied to citrus seedlings grown in greenhouses or shadehouses.			
<b>Restriction:</b>			
<ul style="list-style-type: none"> <li>Minimum retreatment interval is 7 days.</li> </ul>			
<b>CITRUS</b> <b>Field Nursery Grown</b>			
<p>To control Melanose, Scab, Pink Pitting, Greasy Spot, Brown Rot and for suppression of Citrus Canker, apply 2 to 6.3 pounds of Kocide® 50DF per acre. Apply Kocide® 50DF at 28-day intervals if needed depending on disease severity. Minimum retreatment interval is 7 days. Maximum Annual Rate/Acre is 25.1 lbs.</p>			

**Crop-Specific Restrictions:**

If Kocide® 50DF is applied in a spray solution having a pH of less than 6.5, phytotoxicity may occur.

Do not tank mix Kocide® 50DF with any product containing the active ingredient aluminum tris (O-ethyl phosphonate) for use on any registered crops unless appropriate precautions have been taken to buffer the spray solution because severe phytotoxicity may result.

Refer to EPA-approved label for complete list of **USE RESTRICTIONS**.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the precautionary statements plus all other information appearing on the product labels. **This recommendation must be in the possession of the user at the time of pesticide application.**

