

## FIFRA Section 2(ee) Recommendation

For suppression of Root Knot (*Meloidogyne* spp.), Cyst (*Globodera* spp.), Lesion (*Pratylenchus* spp.), Ring (*Mesocriconema* spp.) and Spiral (*Helicotylenchus* spp.) nematodes

**FOR USE AND DISTRIBUTION IN THE STATE OF TEXAS ONLY**

**Expires 12/31/2024**

*This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the US Environmental Protection Agency.*

All applicable directions, restrictions, and precautions on the EPA-registered labels must be followed. Please contact your regional sales manager to determine the specific requirements for FIFRA §2(ee) recommendations in your respective state.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Crops: All labeled crops**

**Application Rate: 1-2 quarts per acre as part of a root disease program for nematode suppression.**

**Root disease soil application:** Mix the required amount in sufficient water to apply by one of the following methods:

- Soil drench at transplanting, using a “water wheel” injector, spray nozzles/hoses, or other method to drench each root ball and/or planting hole.
- Soil or seed line drench, or banded spray (in-furrow) at planting (refer to label for specific instructions).

**Follow-up (post-planting):** Preventative applications can be made every 2-4 weeks as needed by one or more of the following methods:

- Drip (trickle) or any type of sprinkler irrigation, any time after planting or transplanting. See Chemigation Instructions for additional information.
- Directed spray onto the soil surface and/or lower plant parts. If targeting root disease, follow immediately with sufficient overhead sprinkler irrigation to move Double Nickel® LC to the root zone.
- Injection directly into the rooting zone using shanks or similar equipment.

### **Crop-Specific Restrictions:**

Do not apply when wind speed favors drift beyond the area intended for treatment.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the precautionary statements plus all other information appearing on the product labels. **This recommendation must be in the possession of the user at the time of pesticide application.**

Double Nickel® LC 2(ee) Nematodes (TX)

Issued: 2021-11-10

Page 1 of 1



Certis USA LLC  
9145 Guilford Road, Suite 175  
Columbia MD 21046  
(800) 250-5024

[www.certisbio.com](http://www.certisbio.com)