

FIFRA Section 2(ee) Recommendation

For Suppression of Spotted Wing Drosophila (*Drosophila suzukii*) in Berries, including Blackberry, Blueberry, Gooseberry, Grape, Loganberry, Raspberry, and Strawberry.

FOR USE AND DISTRIBUTION IN THE STATE OF TENNESEE ONLY

Expires 06/30/2031

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the US Environmental Protection Agency.

All applicable directions, restrictions, and precautions on the EPA-registered labels must be followed. Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA §2(ee) recommendations in your respective state.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop: Berries including blackberry, blueberry, gooseberry, grape, loganberry, raspberry, and strawberry

Application Rate: Apply 0.25% to 1% solution (1-4 quarts Sil-MATRIX[®] LC/100 gal.) in 50 to 250 gallons finished spray per acre.

Crop-Specific Restrictions:

Do not apply more than 10 quarts per acre (7.5 lb ai/a) per application.

Do not apply more than 20 gallons per acre (60 lb ai/a) per season.

Do not make post-harvest applications.

Refer to EPA-approved label for complete list of **USE RESTRICTIONS**.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the precautionary statements plus all other information appearing on the product labels. **This recommendation must be in the possession of the user at the time of pesticide application.**

